

订购热线: 4008-898-798

Anti-REL antibody

Cat. No. ml122147

Package 25 μl/100 μl/200 μl

Storage -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Product overview

Description Anti-REL rabbit polyclonal antibody

Applications ELISA, WB, IHC

Immunogen Fusion protein of human REL

ReactivityHumanContent0.4 mg/mlHost speciesRabbit

Ig classImmunogen-specific rabbit IgGPurificationAntigen affinity purification

Target information

Symbol REL

Full name v-rel avian reticuloendotheliosis viral oncogene homolog

Synonyms C-Rel Swissprot Q04864

Target Background

The REL gene encodes c-Rel, a transcription factor that is a member of the Rel/NFKB family, which also includes RELA (MIM 164014), RELB (604758), NFKB1 (MIM 164011), and NFKB2 (MIM 164012). These proteins are related through a highly conserved N-terminal region termed the 'Rel domain,' which is responsible for DNA binding, dimerization, nuclear localization, and binding to the NFKB inhibitor




订购热线: 4008-898-798


Applications

Immunohistochemistry

Predicted cell location: Nucleus or Cytoplasm

Positive control: Human brain Recommended dilution: 50-200





The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using ml122147(REL Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: ×200)

Western blotting

Predicted band size:69 kDa Positive control:293T cells

Recommended dilution: 500-2000

Gel: 6%SDS-PAGE


Lysate: 40 µg


Primary antibody: ml122147(REL Antibody) at dilution 1/950

Exposure time: 10 seconds

Predicted cell location: Nucleus or Cytoplasm Positive control: Human prostate cancer

Recommended dilution: 50-200






The image on the left is immunohistochemistry of paraffin-embedded Human prostate cancer tissue using ml122147(REL Antibody) at dilution 1/30, on the right is treated with fusion protein. (Original magnification: ×200)

Lane: 293T cells

eisakii produce Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution



ELISA

Recommended dilution: 2000-5000

联系电话: 4008-898-798, 021-61725725

联系QQ: 2881505695, 2881505696

邮箱: mlbio cn@yeah.net

网址: www.mlbio.cn