

兔抗 DACT2 多克隆抗体

中文名称：兔抗 DACT2 多克隆抗体

英文名称：Anti-DACT2 rabbit polyclonal antibody

别 名：DPR2; DAPPER2; C6orf116; bA503C24.7

相关类别：一抗

储 存：冷冻（-20℃）

宿 主：Rabbit

抗 原：DACT2

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

| | |
|----------------------|---|
| Background: | Involved in regulation of intracellular signaling pathways during development. Negatively regulates the Nodal signaling pathway, possibly by promoting the lysosomal degradation of Nodal receptors, such as TGFBR1. May be involved in control of the morphogenetic behavior of kidney ureteric bud cells by keeping cells epithelial and restraining their mesenchymal character. May play an inhibitory role in the re-epithelialization of skin wounds by attenuating TGF-beta signaling (By similarity). |
| Applications: | ELISA, IHC |

| | |
|------------------------------------|--|
| Name of antibody: | DACT2 |
| Immunogen: | Fusion protein of human DACT2 |
| Full name: | dishevelled binding antagonist of beta catenin 2 |
| Synonyms: | DPR2; DAPPER2; C6orf116; bA503C24.7 |
| SwissProt: | Q5SW24 |
| ELISA Recommended dilution: | 5000-10000 |
| IHC positive control: | Human ovarian cancer; |
| IHC Recommend dilution: | 25-100 |

