

兔抗 CELSR1 多克隆抗体

中文名称：兔抗 CELSR1 多克隆抗体

英文名称：Anti-CELSR1 rabbit polyclonal antibody

别名：cadherin EGF LAG seven-pass G-type receptor 1; ME2; FMI2; CDHF9; HFMI2; ADGRC1

相关类别：一抗

抗原：CELSR1

储存：冷冻（-20℃）

宿主：Rabbit

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

技术规格

Background:

The protein encoded by this gene is a member of the flamingo subfamily, part of the cadherin superfamily. The flamingo subfamily consists of nonclassic-type cadherins; a subpopulation that does not interact with catenins. The flamingo cadherins are located at the plasma membrane and have nine cadherin domains, seven epidermal growth factor-like repeats and two laminin A G-type repeats in their ectodomain. They also have seven transmembrane domains, a characteristic unique to this subfamily. It is postulated that these proteins are receptors involved in contact-m

| | |
|------------------------------------|--|
| | <p>mediated communication, with cadherin domains acting as homophilic binding regions and the EGF-like domains involved in cell adhesion and receptor-ligand interactions. This particular member is a developmentally regulated, neural-specific gene which plays an unspecified role in early embryogenesis.</p> |
| Applications: | ELISA, IHC |
| Name of antibody: | CELSR1 |
| Immunogen: | Synthetic peptide of human CELSR1 |
| Full name: | cadherin EGF LAG seven-pass G-type receptor 1 |
| Synonyms: | ME2; FMI2; CDHF9; HFMI2; ADGRC1 |
| SwissProt: | Q9NYQ6 |
| ELISA Recommended dilution: | 500-1000 |
| IHC positive control: | Human liver cancer and human colorectal cancer |
| IHC Recommend dilution: | 25-100 |



